



HAL
open science

Elastic properties of the zygomatic muscle using ultrasound elastography technique

Redouane Ternifi, Philippe Pouletaut, Adrien Heintz, Stéphanie Dakpé, Sylvie Testelin, Bernard Devauchelle, Fabrice Charleux, Jean-Marc Constans, Sabine Bensamoun

► **To cite this version:**

Redouane Ternifi, Philippe Pouletaut, Adrien Heintz, Stéphanie Dakpé, Sylvie Testelin, et al.. Elastic properties of the zygomatic muscle using ultrasound elastography technique. 8th World Congress of Biomechanics, Jul 2018, Dublin, Ireland. hal-01982826

HAL Id: hal-01982826

<https://hal.utc.fr/hal-01982826>

Submitted on 16 Jan 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Elastic properties of the zygomatic muscle using ultrasound elastography technique

Redouane Ternifi¹, Philippe Pouletaut¹, Adrien Heintz², Stéphanie Dakpé³, Sylvie Testelin³,
Bernard Devauchelle³, Fabrice Charleux⁴, Jean-Marc Constans², Sabine F. Bensamoun¹

¹ Sorbonne University, Université de technologie de Compiègne CNRS, UMR 7338 Biomechanics and Bioengineering, centre de Recherche de Royallieu, Compiègne, France

² CHU Amiens-Institut Faire Faces, Imagerie et Radiologie Médicale, Amiens, France

³ CHU Amiens-Institut Faire Faces, service de chirurgie maxillo-faciale, Amiens, France

⁴ ACRIM-Polyclinique Saint Côme, Compiègne, France

Introduction

The face expression is animated by the movements of the muscles, which can be altered during pathology. The main objective of this study is to develop an experimental imaging protocol to evaluate the functional properties of the zygomaticus major (ZM) muscle using Magnetic Resonance Imaging (MRI) and ultrasound (US) elastography techniques. In addition, the metabolic behavior of the ZM has been characterized with MR Spectroscopy (MRS).

Methods

Eight healthy volunteers (age range: 21 - 30 years) without face muscle damage have been recruited at Amiens Hospital (France). Three capsules full of oil have been placed in three face area (nose, ear and chin), allowing a tracking accuracy for ultrasound fusion process. Subsequently, each subject has undergone MRI, MRS and US elastography tests. 3D-T1 MRI sequence (Philips®, 3T) has been performed with different coils to localize the ZM and the tracking components. MRS has been applied in the center of the ZM using different TE (35ms, 144ms, 288ms and 432ms). Four spectral profiles showing the ZM metabolic behavior have been recorded. After MRI/MRS acquisitions, B-mode ultrasound (SLH20-6 probe), shear wave elastography (SL10-2 probe) and fusion process have been performed in the same ZM area using an Aixplorer® machine. Morphological parameters, elastic properties and superimposition of the ZM anatomical images (MRI & US) have been determined.

Results

The SLH20-6 probe, clinically used for hand tendon, provides a more accurate definition of the face muscle structure compared to the SL10-2 probe. The mean thickness of the ZM is 1.83 ± 0.28 mm. A better signal to noise has been obtained with the dStream Head 32 channels. The mean elastic properties were 19.60 ± 2.63 kPa. These values are in the same range as other head and neck muscles [1]. The fusion process, clinically used for abdominal tests, has been successfully developed for the face demonstrating an accurate superimposition of MRI and US face tissues. MRS acquisitions have identified well known metabolites usually present in muscle tissue [2].

Discussion

The originality of this study has been to provide new functional, morphological and metabolism properties of the zygomaticus major muscle. The present protocol could be applied to evaluate the effects of treatment (such as injection inside the ZM) on the functional properties. These measurements will help clinician to adapt and follow the treatment. Future MRS measurements will be performed to accurately identify the ZM metabolism behavior which could be correlated to the elastic properties.




Figure 1: A) elasticity map obtained with ultrasound elastography technique. B) Spectral profile (TE = 144ms) obtained with MRS technique.

Acknowledgements

This work was supported by the Hauts de France Region.

References

- [1] Dieterich et al. 2016. Eur J Appl Physiol.
- [2] Pola et al. 2012. Progress in Nuclear Magnetic Resonance Spectroscopy.